

Goldsmiths Research Online

*Goldsmiths Research Online (GRO)
is the institutional research repository for
Goldsmiths, University of London*

Citation

Pluess, Michael; De Brito, Stephane A.; Jones Bartoli, Alice; McCrory, Eamon and Viding, Essi. 2022. Individual differences in sensitivity to the early environment as a function of amygdala and hippocampus volumes: An exploratory analysis in 12-year-old boys. *Development and Psychopathology*, 34(3), pp. 901-910. ISSN 0954-5794 [Article]

Persistent URL

<https://research.gold.ac.uk/id/eprint/30295/>

Versions

The version presented here may differ from the published, performed or presented work. Please go to the persistent GRO record above for more information.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Goldsmiths, University of London via the following email address: gro@gold.ac.uk.

The item will be removed from the repository while any claim is being investigated. For more information, please contact the GRO team: gro@gold.ac.uk

Running head: SENSITIVITY & BRAIN STRUCTURE

**Individual Differences in Sensitivity to the Early Environment as a Function of
Amygdala and Hippocampus Volumes: An Exploratory Analysis in 12-Year Old
Boys**

Michael Pluess¹

Stephane A. De Brito²

Alice Jones Bartoli³

Eamon McCrory⁴

Essi Viding⁴

¹ Department of Biological and Experimental Psychology, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom

² School of Psychology, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

³ Department of Psychology, Goldsmiths University of London, Whitehead Building New Cross, London SE14 6NW, United Kingdom

⁴ Psychology and Language Sciences, University College London, 26 Bedford Way London WC1E 7HB, United Kingdom

Acknowledgments

We gratefully acknowledge the ongoing contribution of the participants in TEDS and their families. TEDS is supported by a program grant to Robert Plomin from the UK Medical Research Council [MR/M021475/1; and previously G0901245], with

additional support from the US National Institutes of Health [HD044454; HD059215; AG046903] and the European Commission [602768]. Collection of the neuroimaging data reported here was supported by grant G0401170 from the Medical Research Council and grant MRD 12-73 from the UK Department of Health Forensic Mental Health Programme awarded to Essi Viding. Stephane De Brito was supported by an advanced research fellowship from the Swiss National Science Foundation (FNS PA00P1_139586).

Declaration of Interests

The authors declare no conflict of interest.

Correspondence to:

Michael Pluess, Department of Biological and Experimental Psychology, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom

Email: m.pluess@qmul.ac.uk, Phone: +44 (0)207-882-8004

Individual Differences in Sensitivity to the Early Environment as a Function of Amygdala and Hippocampus Volumes: An Exploratory Analysis in 12-Year Old Boys

Abstract

Children differ in their response to environmental exposures with some being more sensitive to contextual factors than others. According to theory, such variability is the result of individual differences in neurobiological sensitivity to environmental features with some being generally more affected by both negative and/or positive experiences. In this exploratory study we tested whether left and right amygdala and hippocampus volumes (corrected for total brain size) account for individual differences in response to environmental influences in a sample of 62 boys. Cumulative general environmental quality, ranging from low to high, was measured across the first nine years and child behavior was reported by teachers when boys were 12-13 years old. According to analyses, only the left amygdala volume, but not any of the other brain volumes, emerged as an important brain region for sensitivity to *positive* environmental aspects. Boys with a larger left amygdala benefited significantly more from higher environmental quality than boys with a smaller left amygdala whilst not being more vulnerable to lower quality. Besides providing preliminary evidence for differences in *Environmental Sensitivity* due to brain structure, results also point to the left amygdala as having a specific role regarding the response to environmental influences.

Keywords: Differential Susceptibility; Vantage Sensitivity; Environmental Sensitivity; Amygdala; Hippocampus;

**Individual Differences in Sensitivity to the Early Environment as a Function of
Amygdala and Hippocampus Volumes: An Exploratory Analysis in 12-Year Old
Boys**

Like many other species, humans are able to adapt to a wide range of environmental conditions and exposures (Bateson et al., 2004). Such adaptation is crucial for successful development given that different contexts may require different behavioral strategies (e.g., aggression in hostile conditions versus cooperation in supportive contexts) in order to achieve survival and reproduction (Belsky & Pluess, 2009b). However, effective adaptation is conditional on accurate perception and interpretation of environmental cues. The general ability to register and process external stimuli has been defined as *Environmental Sensitivity* (Pluess, 2015). Empirical studies report substantial inter-individual differences in the degree of Environmental Sensitivity as a function of genetic, physiological, and psychological factors (Belsky & Pluess, 2009a, 2011). These sensitivity factors are hypothesized to be markers of a more responsive central nervous system (Pluess, Stevens, & Belsky, 2013). In the current study we test this *neurosensitivity* hypothesis in an exploratory analysis by investigating whether individual differences in brain structure of the amygdala and hippocampus, assessed in early adolescence, moderate the association between broad environmental quality across childhood and behaviour problems in early adolescence (accounting for potential correlations between childhood environment and brain structure).

It can be widely observed that humans, like many other species (Wolf, van Doorn, & Weissing, 2008), differ substantially in their response to various environmental exposures and experiences. Such differences have traditionally been interpreted from a perspective of *Diathesis-Stress* with a focus on individual differences in vulnerability to adversity (Zuckerman, 1999). The *Diathesis-Stress*

perspective suggests that certain people, those characterised by some kind of vulnerability (e.g., genetic or psychological traits), are more likely to develop psychological problems when experiencing adversity. However, the model does not make any predictions about differences in the response to positive experiences. Over the last decade several theories have been put forward suggesting that people differ in their sensitivity to environmental quality, with some being generally more and some generally less sensitive (Belsky & Pluess, 2009a; Boyce & Ellis, 2005; Ellis, Boyce, Belsky, Bakermans-Kranenburg, & Van IJzendoorn, 2011). For example, the frameworks of *Differential Susceptibility* (Belsky & Pluess, 2009a), *Biological Sensitivity to Context* (Boyce & Ellis, 2005), and *Sensory Processing Sensitivity* (Aron & Aron, 1997) all propose that individuals with higher sensitivity are not just more reactive to negative experiences but also more sensitive to positive environmental influences, compared to less sensitive people. The notion that people differ in their response to *positive* experiences with some benefitting disproportionately more than others has recently been formalized more specifically in the framework of *Vantage Sensitivity* and is supported by a growing number of empirical studies (Pluess & Belsky, 2013) (see Figure 1 for a graphical illustration of *Diathesis-Stress*, *Vantage Sensitivity*, and *Differential Susceptibility*).

--- Insert Figure 1 here ---

Drawing on evolutionary considerations, both *Differential Susceptibility* (Belsky & Pluess, 2009a) and *Biological Sensitivity to Context* (Boyce & Ellis, 2005) propose that natural selection shaped individual differences in the propensity for conditional adaptation to the environment because both low and high sensitive

phenotypes have particular evolutionary benefits, depending on the specific quality of the developmental context (Ellis et al., 2011). For example, while low sensitive individuals tend to be more resilient in the face of adversity, they also suffer the disadvantage of being less responsive to supportive aspects of the environment. Highly sensitive individuals, on the other hand, are more likely to benefit from positive environmental exposures whilst also being more vulnerable to the negative effects of adverse experiences (Pluess, 2015). As a consequence, the general population is made up of individuals that differ in their general environmental sensitivity with a significant minority of about 30% characterised by particularly high sensitivity (Lionetti et al., 2018; Pluess et al., 2018).

A large number of studies provide empirical evidence that some people are more sensitive than others to both negative *and* positive environmental influences (Belsky & Pluess, 2016; Slagt, Dubas, Dekovic, & van Aken, 2016). For example, sensitive children (i.e., those with a high-reactive temperament in infancy) have been shown to develop more behavioural problems than less sensitive children when experiencing harsh parenting but also as having fewer problems when parenting was especially sensitive (Slagt et al., 2016). Similarly, children carrying more sensitivity genes (based on a recently developed genome-wide polygenic sensitivity score) presented with more emotional problems than genetically less sensitive children when experiencing negative parenting but also significantly fewer problems than other children when exposed to positive parenting (Keers et al., 2016).

Although sensitivity markers have been identified across different levels of analysis including the genetic, physiological and behavioral level, the exact mechanism(s) underlying individual differences in Environmental Sensitivity is not yet well understood. However, all leading theories in the field converge on the

hypothesis that features of the central nervous system play a central role (Acevedo et al., 2014; Aron, Aron, & Jagiellowicz, 2012; Ellis et al., 2011). This view is supported by the notion that Environmental Sensitivity requires a vast array of highly complex cognitive functions, including sensory perception, processing of sensory intake, emotional reactivity as well as higher order cognitive functions such as attention, memory, and executive function. Hence, the various sensitivity markers are understood to contribute to—or manifest—features of a more sensitive central nervous system (Aron et al., 2012; Ellis et al., 2011; Pluess, 2015). While this *neurosensitivity* hypothesis (Pluess & Belsky, 2013; Pluess et al., 2013) is rather broad and unspecific, studies reporting associations between established sensitivity markers and both structural (e.g., Holmes et al., 2012) and functional measures of the brain (e.g., Munafo, Brown, & Hariri, 2008; Schwartz, Wright, Shin, Kagan, & Rauch, 2003), point to the amygdala and hippocampus as key regions of interest (although other brain regions and networks likely play a role too). For example, several candidate genes that have been associated with heightened sensitivity to the environment (Belsky et al., 2009; Belsky & Pluess, 2009a, 2013) have been linked to amygdala or hippocampus function or structure, such as genetic variation in the serotonin transporter gene (5-HTTLPR; e.g., Hariri et al., 2002), COMT (e.g., Rasch et al., 2010), and MAOA (e.g., Meyer-Lindenberg et al., 2006). Similarly, behavioral markers of sensitivity, including high reactivity in infancy (Schwartz et al., 2012), inhibited temperament at five years (Hill, Tessner, Wang, Carter, & McDermott, 2010), and behavioral inhibition in adults (Barros-Loiscertales et al., 2006) have been associated with amygdala function and structure.

In the current paper we focus on structural (rather than functional) differences in these two regions, given that a small number of studies provided first evidence that

structural differences in these brain regions indeed moderate the effects of environmental influences consistent with what *Environmental Sensitivity* frameworks predict. For example, one cross-sectional study featuring a sample of 106 11-14 year old adolescents and their mothers, aimed at investigating whether structural differences in the amygdala, hippocampus, and anterior cingulate cortex moderated the relationship between observationally assessed maternal aggression and adolescents' depression symptoms (Yap et al., 2008). Several statistically significant three-way interactions between maternal aggression, brain structure, and gender emerged. Most relevant for the current paper, girls with *smaller* (bilateral) amygdala volumes were more sensitive to maternal aggression *for better and for worse*. They had the lowest depression scores when maternal aggression was low and the highest depression scores when mothers were more aggressive, compared to girls with larger amygdala. However, in boys a *larger* right amygdala was associated with reduced depression symptoms when maternal aggression was low (i.e., reflecting *Vantage Sensitivity*) with no evidence for heightened vulnerability to high maternal aggression. In a follow-up study of the same sample, the authors tested whether individual differences in hippocampal volume moderated change in depression scores over time (Whittle et al., 2011). Again, a three-way interaction emerged but this time suggesting that larger (bilateral) hippocampal volume predicted sensitivity to maternal aggression in the prediction of depression, but only in girls. Those with larger hippocampal volumes had the lowest depression scores when maternal aggressive behavior was low, and the highest depression scores when maternal aggressive behavior was high. Depression symptoms in girls with smaller hippocampal volumes, on the other hand, were less affected by maternal aggression. More recently, a longitudinal study involving 209 17-year old adolescents tested whether hippocampal volume moderated

the effects of family connectedness and community crime on depression, anxiety, and externalizing symptoms (Schriber et al., 2017). The effects of both environmental measures on depression were moderated by the volume of the left hippocampus, across both genders. According to follow-up analyses, adolescents with a larger left hippocampus had higher depression scores when family connectedness was low and when community crime was high, consistent with *Diathesis-Stress*.

In summary, these three studies provide important first evidence that individual differences in Environmental Sensitivity are associated with differences in amygdalar and hippocampal volumes. However, this pioneering research was primarily informed by a pathology framework focused on vulnerability to the negative effects of specific adverse experiences. Consequently, most of the studies did not consider whether brain structure also predicted individual differences at the positive end of the environmental quality spectrum. Furthermore, outcome measures were exclusively based on self-report and significant interactions were not followed up with the statistical procedures required to explore whether detected interaction patterns are more consistent with *Diathesis-Stress*, *Differential Susceptibility*, or *Vantage Sensitivity* (Roisman et al., 2012).

The current analysis builds on the studies mentioned above, but rather than focusing on specific childhood experiences, it features a more general and cumulative measure of broad environmental quality across childhood, ranging from low (i.e., negative) to high (i.e., positive). Such a broad and inclusive measure of environmental quality may be more suitable when investigating individual differences in general Environmental Sensitivity, as is the case in the current study. In addition, our study examines the pattern of significant interactions to test whether interactions are more supportive of *Diathesis-Stress*, *Differential Susceptibility*, or *Vantage Sensitivity*.

Importantly, the study is exploratory in nature given that it is based on data from a relatively small number of children, initially recruited for a study with different aims, and including only boys. The main objective of this proof of concept study was to investigate whether differences in amygdalar and hippocampal volumes, assessed in early adolescence, moderate the effects of a cumulative score of broad early environmental quality, on teacher-reported child behavior (i.e., an index of emotional and behavioral problems as well as prosocial skills) in early adolescence.

Methods

Participants

Rather than recruiting new participants, we relied on a data set that was already available to us. Important to mention, the available sample did not include any girls and was made up of male twins. Data was obtained from 62 boys originally included in the *Twins Early Development Study* (TEDS), a large longitudinal study of over 16,000 twin pairs born in England and Wales in 1994, 1995, and 1996. TEDS includes extensive data on various aspects of development, collected at regular intervals from a sample that is representative of the UK population (Kovas et al., 2007). General data and recruitment procedures for TEDS are reported in detail elsewhere (Haworth, Davis, & Plomin, 2013). The participants of the current study are a subset of boys that were initially recruited from the TEDS database for an associated twin neuroimaging project on precursors of psychopathic traits led by authors of the current study (Rijsdijk et al., 2010). Parents completed questionnaires about MRI contra-indicators and provided consent to be contacted regarding study participation. After description of the study to the children and their parents, written informed consent was obtained from parents and oral assent from boys. Study and recruitment procedure were approved by the Institute of Psychiatry and South London

and Maudsley NHS Foundation Trust Research Ethics Committee. For the current analysis only boys with complete data on all measures and very good quality imaging data (i.e., no blurring by motion) were included, resulting in a total sample of $N = 62$ boys with a mean age of 11.61 years ($SD = .81$) at the time of imaging (see Table 1 for a detailed description of the sample). The participating boys were aged 10–13 years, had no psychiatric, neurological or medical problems, and a full scale IQ of at least 80. Premature and low birthweight children were excluded from analyses. All boys included in the current analysis come from same sex twin pairs (20 MZ twin siblings, 42 DZ twin siblings). Relatedness between boys from the same twin pair (the sample included 2 complete MZ twin pairs and 16 complete DZ twin pairs) was statistically controlled for by nesting related siblings in hierarchical linear models. Compared to the full TEDS sample, the boys included in the current study were significantly older when behavior problems were rated by teachers ($M = 12.09$, $SD = .29$, for included cases versus $M = 11.53$, $SD = .66$ for the remaining TEDS sample, with $t = -14.78$, $p < .01$). Furthermore, total problem scores were slightly higher in this subsample of boys compared to the complete TEDS sample ($M = 6.50$, $SD = 5.57$, for included cases versus $M = 5.13$, $SD = 5.16$ for other TEDS cases with $t = -2.08$, $p = .04$) which can be explained by the fact that a proportion of the sample recruited for the original neuroimaging project on psychopathic traits was selected based on elevated conduct problems and callous-unemotional scores (i.e., those scoring in the top 10% on these measures) when they were nine years old. Importantly, 42 of the boys included in the current analysis represent control cases with normal levels of behavioral problems (i.e., the current sample does not reflect high risk for psychopathology). Control cases had significantly lower behavioral problems ($M = 5.21$, $SD = 4.9$) than those selected for elevated problems ($M = 9.04$, $SD = 6.04$; with t

= 2.69, $p = .01$). However, boys with elevated problems did not differ from controls regarding amygdala or hippocampus volumes or any other variable included in the analysis. Furthermore, the cumulative environmental quality score of the subsample did not differ from the total sample ($M = 13.88$, $SD = 1.67$, of included cases versus $M = 13.69$, $SD = 1.89$ of other TEDS participants with $t = -.80$, $p = .42$). According to chi-square tests, there were no significant differences in ethnicity and income between the current sample and the remaining TEDS sample.

MRI Acquisition

Structural brain images were acquired using a General Electric Signa 3.0 Telsa Excite II MRI scanner (GE Medical systems, Milwaukee, WI, USA) at the Centre for Neuroimaging Science, Institute of Psychiatry, London, UK. A high-resolution, 3D T1 weighted dataset was acquired using an inversion recovery prepared spoiled gradient echo (IR-SPGR) sequence. Imaging parameters were TR = 8ms; TE = 2.9ms; TI = 450ms; excitation flip angle = 20°. The in-plane matrix size was 256x192 over a 280x210mm field of view, reconstructed to 256x256 over 280x280mm. In plane pixel size was thus 1.094 x 1.094 mm. Two hundred through plane partitions (each 1.1mm thick) were collected, with two partitions being discarded at each end of the imaging volume to minimize wrap-round artefacts. Partial k-space coverage ('0.75 NEX') was used. The scanning time was 6 minutes.

MRI data pre-processing

To quantify and extract the grey matter volume from the amygdala and the hippocampus, we used voxel-based morphometry (VBM), a widely employed automated MRI analysis technique (Whitwell, 2009). Specifically, the data were pre-processed using the VBM8 toolbox and SPM8, which provide improved segmentation and registration procedures such as the Diffeomorphic Anatomical Registration

Through Exponentiated Lie Algebra (DARTEL) toolbox (Ashburner, 2007). Given that our sample included children aged 10-13 years, customised tissue probability maps were created in the Montreal Neurological Institute (MNI) space for use with the VBM8 Toolbox. These were produced using the matched template approach of the Template-O-Matic Toolbox for SPM8 with each participant's age and sex as defining variables (Wilke, Holland, Altaye, & Gaser, 2008). The pre-processing included the following steps: First, the anterior commissure was manually indicated on all structural images as the [0, 0, 0 mm] origin in the MNI spatial coordinate system. Individual images were then corrected for bias-field inhomogeneities, segmented and spatially normalised (affine-only transformation) with reference to customised tissue probability maps. Segmentation accuracy was visually checked for each participant. Based on individual registered grey matter and white matter segmentations, an average DARTEL template of all participants was created in MNI space (Ashburner, 2007). The affine-registered grey matter and white matter segments were then warped to this average template using the high-dimensional DARTEL approach. Importantly, in order to account for individual differences in brain size modulated data produced gray matter volume (GMV) and the voxel values in the grey matter segments were only multiplied by the non-linear component of the registration. Given that the two structures we focused on are well-defined and adjacent to each other, we did not smooth the data. For each participant, the MarsBaR region of interest (ROI) toolbox (Brett et al., 2002; Tzourio-Mazoyer et al., 2002) as implemented in SPM8 was used to extract mean grey matter volume value for the amygdala and the hippocampus, bilaterally, using the anatomical masks from the aal atlas (Figure 2).

--- Insert Figure 2 here ---

Measures

Environmental Quality. In order to obtain a measure of general and broad environmental quality across childhood, a new scale for cumulative environmental quality across years 1-9 was created by recoding and summing up existing parent-reported variables and scales reflecting both negative and positive aspects of the social and material developmental context. All included variables reflect normative aspects of children's environment rather than unusual experiences such as severe adversity. The subscale *Social Environment* covers six parent-reported psycho-social aspects (i.e., parental feelings, parental discipline, family order and chaos, partner presence, parental reading to child, parent outings with child, all assessed at 3 years and then again at 9). The subscale *Material Environment* is based on six parent-reported socio-economic characteristics of the environment (i.e., socio-economic status composite scores at 1st contact and at 7 years, number of books at 3 years, as well as financial changes, parental unemployment, and household income at 9 years). After recoding all individual variables to fit a categorical scale with "1 = low quality", "2 = medium quality", and "3 = high quality", items were summed up separately for each of the subscales (see supplementary information for details). Importantly, according to preliminary analyses using the complete TEDS sample, the recoded items were all significantly and negatively associated with child total problems (SDQ) rated by parents (i.e., the higher the quality, the lower the problems). The separate sum scores for the social environment at ages 3 and 9 years were averaged to compute a *Social Environment* subscale that reflects environmental quality across childhood. In order to create the final *Environmental Quality* score, the *Social Environment* subscale score was then averaged with the *Material Environment* subscale score. This

resulted in a total score with a theoretical range of 8-18 with 8 = lowest quality and 18 = highest quality.

Child Behavior. The Strengths and Difficulties Questionnaire (SDQ), a well-established 25-item questionnaire with five subscales (Goodman, 1997), was completed by teachers when children were 12-13 years old. Items were rated on a Likert scale ranging from “0 = not true“, “1 = somewhat true”, to “2 = certainly true”. The subscales (i.e., emotional symptoms, conduct problems, hyperactivity/inattention, peer relationship problems and prosocial behavior) can be used individually (each ranging from 0-10) but the four problem subscales can also be combined to yield a total problem score (with a range of 0-40). For the current analysis we focused on the total problem score which included the subscales *emotional symptoms, conduct problems, hyperactivity/inattention, and peer relationship problems*. In a second step we also conducted a set of post-hoc analyses for each individual subscale of the SDQ (see supplementary information).

Statistical Analysis

Interactions between brain volumes and environmental quality predicting total behavior problems were tested with hierarchical linear models. Relatedness between siblings was controlled for by including the family level as a random effect (i.e., data of related siblings were nested in families). Models were kept as parsimonious as possible given the small sample size and tested for the main effects of the environmental quality score and continuous brain volumes as well as the interaction between both. Given we had no hypothesis to assume sensitivity would be different as a function of zygosity (according to bivariate correlations zygosity was unrelated to all included variables, $ps > .05$), we did not control for MZ or DZ status. Importantly, all brain volumes measures were corrected for total brain size (modulation with non-

linear registration; see MRI pre-processing section) and age at the time of imaging was not significantly correlated with any of the volumes ($r = -.10$ to $.12$ with $p = .25-.90$). Separate models were run for the four different brain regions (i.e., left and right amygdala, left and right hippocampus). In order to follow-up significant interactions the continuous brain volume variable was divided into low and high volume at the median to create and compare simple slopes. Finally, regions of significance analyses were conducted in order to examine whether detected interactions were more supportive of *Diathesis-Stress*, *Vantage Sensitivity*, or *Differential Susceptibility* patterns (Preacher, Curran, & Bauer, 2006). In addition, significant interactions were further investigated in post-hoc analyses that considered the five different subscales of the SDQ separately in relation to *Environmental Quality* but also the subscales of *Social* and *Material Environment*. The level of significance was set at $\alpha = .05$ for all tests but Bonferroni correction that accounts for the correlation between the brain volumes (on average $r = .58$) was applied to correct for multiple testing in relation to the four tested multilevel models. All statistical analyses were conducted in SPSS (version 22).

--- Insert Table 1 here ---

Results

Bivariate Correlations. According to bivariate correlations all four brain volumes were significantly associated with each other (see Table 2). Importantly, brain volumes were not associated with the environmental quality score ($r = -.13$ to $.08$, $p > .05$) or the total problems outcome variable ($r = -.21$ to $.12$, $p > .05$). Finally,

although higher environmental quality was associated with fewer problems, this association did not reach statistical significance ($r = -.14, p > .05$).

--- Insert Table 2 here ---

Hierarchical Linear Models. Interactions between continuous brain volumes and the environmental score were tested with separate hierarchical linear models with biologically related boys nested in families to account for relatedness (i.e., level 1) and controlling for main effects of the cumulative environmental quality score and brain volumes. A significant interaction emerged between left amygdala volume and environmental quality in the prediction of total problems ($B = -16.10, p = .01$). None of the other brain volumes significantly moderated effects of the cumulative environmental score (i.e., interaction terms were $B = -12.32, p = .24$ for right amygdala, $B = 3.48, p = .11$ for left hippocampus, and $B = -11.73, p = .36$ for right hippocampus volume). Importantly, the significant interaction between left amygdala volume and environmental quality survived Bonferroni correction for multiple testing (corrected $\alpha = .028$ taking correlations between the four tested brain volumes into account). The hierarchical linear model results are displayed in Table 3.

--- Insert Table 3 here ---

Follow-up Analyses. The significant interaction was followed-up in order to interpret the nature of the interaction through simple slopes by dividing the sample by median split into two groups, low and high left amygdala volume. In addition to testing simple slopes between environmental score and outcomes for the low and high

left amygdala volume groups we also conducted regions of significance analyses in order to examine the specific pattern of the interaction (Preacher et al., 2006; Roisman et al., 2012). According to simple slopes and regions of significance analysis larger left amygdala volume was associated with higher sensitivity to higher quality of the environment. While the lower bound of the regions of significance analysis was outside of the observed data, the upper bound was within the upper range suggesting that the association between larger left amygdala volume and fewer behavioral outcomes reached statistical significance only towards the higher end of environmental quality. More clearly, whereas environmental quality was not associated with total problems in boys with a small amygdala ($\beta = .16, p = .39$), higher environmental quality was associated with fewer problems in those with a large amygdala ($\beta = -.44, p = .01$), with differences between groups being significant above a threshold of 13.79 on the cumulative environmental quality scale (see Figure 3).

--- Insert Figure 3 here ---

Post-Hoc Analyses. In a series of exploratory post-hoc analyses (not controlling for multiple testing), the hierarchical linear models involving amygdala left volume were rerun separately for the environmental quality score as well as the social environment and material environment subscales predicting all five subscales of the SDQ. Six additional significant interactions emerged: amygdala left volume moderated the effects of the cumulative environmental score on peer ($B = -6.05, p = .01$) and conduct problems ($B = -3.73, p = .02$), the effects of the social environment subscale on emotional symptoms ($B = -3.65, p = .03$) and peer problems ($B = -4.41, p$

= .02), and the effects of the material environment subscale on prosocial behavior ($B = 3.18, p = .05$) and conduct problems ($B = -2.43, p = .02$). The majority of these interactions yielded similar simple slope patterns with larger left amygdala volume reflecting higher sensitivity to higher quality of the early environment. Importantly, interactions effects were not limited to problem behaviors but left amygdala volume also moderated the positive effect of environmental quality on prosocial behavior, a positive outcome, with larger left amygdala being associated with greater prosocial behavior when material environment was particularly high (see Supplementary Information for more information).

Discussion

This exploratory proof of concept study aimed to test whether differences in amygdala and hippocampus volumes moderate the relationship between broad cumulative environmental quality, ranging from low to high, across early childhood and teacher-reported behavioral problems as well as prosocial behavior in early adolescence. Significant interactions were followed-up in order to examine whether the detected sensitivity patterns were more consistent with *Diathesis-Stress* (Zuckerman, 1999), *Differential Susceptibility* (Belsky & Pluess, 2009a), or *Vantage Sensitivity* (Pluess & Belsky, 2013).

In general, results support a *Vantage Sensitivity* pattern for boys with larger amygdala volumes. In more detail, our findings indicate that left amygdala volume moderate associations between early environmental quality and teacher-reported total problems in early adolescence. In boys with a larger left amygdala there was a significant association between higher quality of the environment and lower total problems. This association was not statistically significant in boys with smaller amygdala volumes. The finding that larger left amygdala size in boys reflects higher

sensitivity to higher environmental quality is consistent with a previous study with different predictor and outcome measures, which reported that boys with larger amygdala had fewer depression symptoms when maternal aggression was low (Yap et al., 2008). Interestingly, several studies that investigated the moderating effects of hippocampal volumes (Whittle et al., 2011; Yap et al., 2008) detected such effects in girls, but failed to do so in boys. This appears to be somewhat consistent with observations of the current study that hippocampal volumes did not moderate associations between child behavior and the environment in boys (but one study found that hippocampal volume moderated effects of the social environment across both genders, Schriber et al., 2017). Further studies are required to investigate the potential existence of sex and gender differences regarding associations between brain structure and environmental sensitivity. This is important given well-known sex-specific differences in brain structure, including the amygdala which tends to be generally larger in males (Ruigrok et al., 2014)

Furthermore, according to post-hoc analyses, the moderating effect of left amygdala volume was not restricted to associations between higher environmental quality and lower *maladaptive* outcomes, such as peer problems and emotional problems, but also to higher levels of an *adaptive* outcome, as prosocial behavior. This further corroborates the notion that individuals may differ in their *general* sensitivity to environmental influences rather than exclusively in their vulnerability to develop problem behaviors when exposed to harsh environments (Belsky & Pluess, 2009a). The finding that *larger* amygdala volume in boys growing up in *supportive* environments was not only associated with *fewer* problem behaviors but also with *more* prosocial behavior, is consistent with human and animal studies pointing to associations between amygdala volume and social behavior (Bickart, Wright, Dautoff,

Dickerson, & Barrett, 2011; Dunbar, 2012). Interestingly, follow-up analyses suggest that the significant interaction between amygdala left volume and environmental quality reflects a *Vantage Sensitivity* (Pluess & Belsky, 2013) rather than *Diathesis-Stress* or *Differential Susceptibility* pattern. In other words, boys with larger amygdala volumes were particularly sensitive to higher environmental quality in that they presented with *fewer* problems (and also with more prosocial behaviors) than boys with smaller amygdala volumes when raised in more supportive and more affluent families. In lower quality environments, on the other hand, left amygdala volume was not associated with behavior problems (or prosocial behavior). This is similar to a previous study in which larger amygdala volume of boys predicted *Vantage Sensitivity* to low maternal aggression but not vulnerability to high maternal aggression (Yap et al., 2008). Consequently, this suggests that larger amygdala volume may be an advantage in more supportive and less threatening environments whilst not necessarily increasing vulnerability to adversity—at least in boys that grow up in relatively mild adversity. Future research will have to investigate whether current findings replicate in samples of more deprived backgrounds than the one included in the current study.

Our findings further suggest that structural differences in the amygdala of boys predict the sensitivity to broad and cumulative environmental quality in childhood. Although it is important to identify the specific environmental factors that shape the development of more sensitive children at specific times, combining a large number of environmental aspects across childhood into general and cumulative scores that range from low to high quality and cover various aspects of the normative developmental context appears to be a promising approach when investigating individual differences in more general *Environmental Sensitivity*. Future research

studies may want to adopt this novel approach in order consider sensitivity across a broader spectrum of environmental quality.

The amygdala has been hypothesized as a key region of interest for *Environmental Sensitivity* (Pluess, 2015; Pluess & Belsky, 2013; Pluess et al., 2013) due to associations with various established sensitivity markers. According to empirical studies, larger amygdala volume predicts heightened sensitivity to punishment (Barros-Loscertales et al., 2006) as well as negative affect (Holmes et al., 2012). Importantly, meta-analytic evidence provides strong support for the notion that the amygdala responds to both negative *and* positive stimuli (Sergerie, Chochol, & Armony, 2008). Our research provides further evidence for the amygdala as a brain region of particular relevance for *Environmental Sensitivity*. Although our findings need to be replicated before investigating the specific biological mechanisms underlying the detected associations between amygdala structure and sensitivity in more detail (including potential laterality effects), we suggest that there are good reasons beyond the ones already mentioned why the amygdala might be relevant for sensitivity to the environment. For example, the amygdala's general role in emotional reactivity and response to both negative *and* positive stimuli (Sergerie et al., 2008) may explain why boys with a larger amygdala are more positively affected by higher environmental quality. Furthermore, one of the primary functions of amygdala processing is to signal what is important in any particular situation (i.e., salience processing), and then modulate the appropriate perceptual, attentional, autonomic, and cognitive/conceptual processes to deal with the challenges or opportunities that are present (Cunningham, Arbuckle, Jahn, Mowrer, & Abduljalil, 2010). Children whose brains are more able to do so (e.g., by having a larger amygdala), might adjust better to the conditions of their specific developmental context, such as showing fewer

behavior problems and being more prosocial in a supportive environment.

Importantly, amygdala volume itself has been shown to be associated with environmental conditions in early development such as maternal cortisol during pregnancy (Buss et al., 2012). In light of studies reporting that prenatal and early postnatal factors influence the development of Environmental Sensitivity (Hartman, Freeman, Bales, & Belsky, 2018; Pluess & Belsky, 2011), future studies should also investigate how conditions of the early environment shape characteristics of the brain that are associated with sensitivity to the environment. Moreover, future research should also investigate the role of amygdala *function* (e.g., Gard, Shaw, Forbes, & Hyde, 2018), taking into account recently identified methodological challenges for the use of functional data when researching individual differences (Elliott et al., 2020).

The current study has several important strengths, including prospective longitudinal data, the use of an innovative cumulative environmental score based on objective or parent-reported measures that range across a broad spectrum of environmental quality, as well as teacher-reported outcomes. However, findings have to be considered in light of several limitations. First, the available sample was small and included only twin boys, a third of which had elevated conduct problems and/or callous-unemotional behavior scores at age nine. Future studies should feature larger samples with equal numbers of (non-twin) boys and girls from a general non-clinical population that represents individuals from both low and high socio-economic backgrounds (in order to test whether findings are similar in both low and high risk contexts). In addition, future studies should also investigate whether the findings reported here in early adolescence, extend to other developmental periods across the life span. Second, the sample included some siblings from the same twin pairs which tend to differ less from each other due to shared genetic and environmental factors

(however, we statistically controlled for relatedness by testing hypotheses in a multilevel model and including family as one level within which related individuals were nested in). Third, brain structure was assessed after the environmental exposure occurred and only shortly before teachers rated children's behavior. Ideally, the imaging data would have been obtained before the exposure to exclude the possibility that brain structure itself has been influenced by environmental quality (Evans et al., 2016). Furthermore, although environmental quality was not associated with any of the brain variables according to bivariate correlations, and we also statistically accounted for main effects of brain structure in the model, we cannot exclude the possibility that other environmental factors in early childhood, not assessed with the current measure, contributed to differences in amygdala and hippocampus volumes. Fourth, whilst we used an innovative cumulative environmental quality score in order to capture the general quality of early childhood, this particular score or approach has not been validated through replication in other samples yet. In light of our findings, future studies may want to focus more specifically on measures of environmental enrichment (e.g., frequency of attending museums, shows, exhibitions, after school activities etc.). Fifth, the study did not include important covariates such as gestational age at birth and other pre- and perinatal factors that might influence brain structure (e.g., maternal smoking). Future studies should consider such covariates. Sixth, the current study did not include any explicit measures of sensitivity, such as the *Highly Sensitive Child* scale (Pluess et al., 2018). Future studies should investigate whether validated psychological measures of sensitivity correlate with brain structures found to moderate environmental quality. Finally, the current study only considered amygdala and hippocampus although it is very likely that other brain regions and

networks are also relevant (Acevedo et al., 2014; Moore & Depue, 2016), which should be considered in future studies on sensitivity.

In conclusion, this exploratory study provides novel but preliminary empirical evidence for heightened sensitivity to positive environmental influences for boys in early childhood as a function of structural brain differences in the left amygdala (but not hippocampus) and supports the hypothesis that individual differences in *Environmental Sensitivity* are associated with neurological factors. However, we recommend that our findings be replicated in larger samples of both boys and girls (from high and low socio-economic background), considering a broader range of behavioral outcomes, and with brain structure measured before the environmental exposure.

References

- Acevedo, B. P., Aron, E. N., Aron, A., Sangster, M. D., Collins, N., & Brown, L. L. (2014). The highly sensitive brain: an fMRI study of sensory processing sensitivity and response to others' emotions. *Brain and behavior*, *4*(4), 580-594.
- Aron, E. N., & Aron, A. (1997). Sensory-processing sensitivity and its relation to introversion and emotionality. *Journal of Personality and Social Psychology*, *73*(2), 345-368. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/9248053>
- Aron, E. N., Aron, A., & Jagiellowicz, J. (2012). Sensory processing sensitivity: a review in the light of the evolution of biological responsivity. *Personality and Social Psychology Review*, *16*(3), 262-282. doi:10.1177/1088868311434213
- Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *Neuroimage*, *38*(1), 95-113.
- Barros-Loscertales, A., Meseguer, V., Sanjuan, A., Belloch, V., Parcet, M., Torrubia, R., & Avila, C. (2006). Behavioral inhibition system activity is associated with increased amygdala and hippocampal gray matter volume: a voxel-based morphometry study. *Neuroimage*, *33*(3), 1011-1015.
- Bateson, P., Barker, D., Clutton-Brock, T., Deb, D., D'Udine, B., Foley, R. A., . . . Sultan, S. E. (2004). Developmental plasticity and human health. *Nature*, *430*(6998), 419-421.
- Belsky, J., Jonassaint, C., Pluess, M., Stanton, M., Brummett, B., & Williams, R. (2009). Vulnerability genes or plasticity genes? *Molecular Psychiatry*, *14*(8), 746-754. doi:10.1038/mp.2009.44

- Belsky, J., & Pluess, M. (2009a). Beyond diathesis stress: differential susceptibility to environmental influences. *Psychological Bulletin*, *135*(6), 885-908.
doi:10.1037/a0017376
- Belsky, J., & Pluess, M. (2009b). The Nature (and Nurture?) of Plasticity in Early Human Development. *Perspectives on Psychological Science*, *4*(4), 345-351.
- Belsky, J., & Pluess, M. (2011). Beyond Adversity, Vulnerability and Resilience: Individual Differences in Developmental Plasticity. In D. Cicchetti & G. I. Roisman (Eds.), *Minnesota Symposium on Child Psychology, Vol.36: The Origins and Organization of Adaptation and Maladaptation* (pp. 379-422). Hoboken, NJ: John Wiley.
- Belsky, J., & Pluess, M. (2013). Beyond risk, resilience, and dysregulation: phenotypic plasticity and human development. *Development and Psychopathology*, *25*(4 Pt 2), 1243-1261. doi:10.1017/S095457941300059X
- Belsky, J., & Pluess, M. (2016). Differential Susceptibility to Environmental Influences. In D. Cicchetti (Ed.), *Developmental Psychopathology* (3rd ed., Vol. 3, pp. 59). New York: Wiley.
- Bickart, K. C., Wright, C. I., Dautoff, R. J., Dickerson, B. C., & Barrett, L. F. (2011). Amygdala volume and social network size in humans. *Nature Neuroscience*, *14*(2), 163-164. doi:10.1038/nn.2724
- Boyce, W. T., & Ellis, B. J. (2005). Biological sensitivity to context: I. An evolutionary-developmental theory of the origins and functions of stress reactivity. *Development and Psychopathology*, *17*(2), 271-301. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/16761546>
- Buss, C., Davis, E. P., Shahbaba, B., Pruessner, J. C., Head, K., & Sandman, C. A. (2012). Maternal cortisol over the course of pregnancy and subsequent child

amygdala and hippocampus volumes and affective problems. *Proceedings of the National Academy of Sciences*, *109*(20), E1312-E1319.

Cunningham, W. A., Arbuckle, N. L., Jahn, A., Mowrer, S. M., & Abduljalil, A. M.

(2010). Aspects of neuroticism and the amygdala: chronic tuning from motivational styles. *Neuropsychologia*, *48*(12), 3399-3404.

doi:10.1016/j.neuropsychologia.2010.06.026

Dunbar, R. I. (2012). The social brain meets neuroimaging. *Trends Cogn Sci*, *16*(2),

101-102. doi:10.1016/j.tics.2011.11.013

Elliott, M. L., Knodt, A. R., Ireland, D., Morris, M. L., Poulton, R., Ramrakha, S., . . .

Hariri, A. R. (2020). What Is the Test-Retest Reliability of Common Task-Functional MRI Measures? New Empirical Evidence and a Meta-Analysis.

Psychological Science, 0956797620916786.

Ellis, B. J., Boyce, W. T., Belsky, J., Bakermans-Kranenburg, M. J., & Van

IJzendoorn, M. H. (2011). Differential susceptibility to the environment: An evolutionary–neurodevelopmental theory. *Development and Psychopathology*,

23(01), 7-28. doi:10.1017/S0954579410000611

Evans, G. W., Swain, J. E., King, A. P., Wang, X., Javanbakht, A., Ho, S. S., . . .

Liberzon, I. (2016). Childhood Cumulative Risk Exposure and Adult

Amygdala Volume and Function. *Journal of Neuroscience Research*, *94*(6),

535-543. doi:10.1002/jnr.23681

Gard, A. M., Shaw, D. S., Forbes, E. E., & Hyde, L. W. (2018). Amygdala Reactivity

as a Marker of Differential Susceptibility to Socioeconomic Resources During Early Adulthood. *Developmental Psychology*, *54*(12), 2341-2355.

doi:10.1037/dev0000600

- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38, 581-586.
- Hariri, A. R., Mattay, V. S., Tessitore, A., Kolachana, B., Fera, F., Goldman, D., . . . Weinberger, D. R. (2002). Serotonin transporter genetic variation and the response of the human amygdala. *Science*, 297(5580), 400-403.
- Hartman, S., Freeman, S. M., Bales, K. L., & Belsky, J. (2018). Prenatal stress as a risk—and an opportunity—factor. *Psychological Science*, 29(4), 572-580.
- Haworth, C. M., Davis, O. S., & Plomin, R. (2013). Twins Early Development Study (TEDS): A genetically sensitive investigation of cognitive and behavioral development from childhood to young adulthood. *Twin Research and Human Genetics*, 16(01), 117-125.
- Hill, S. Y., Tessner, K., Wang, S., Carter, H., & McDermott, M. (2010). Temperament at 5 years of age predicts amygdala and orbitofrontal volume in the right hemisphere in adolescence. *Psychiatry Research: Neuroimaging*, 182(1), 14-21.
- Holmes, A. J., Lee, P. H., Hollinshead, M. O., Bakst, L., Roffman, J. L., Smoller, J. W., & Buckner, R. L. (2012). Individual differences in amygdala-medial prefrontal anatomy link negative affect, impaired social functioning, and polygenic depression risk. *Journal of Neuroscience*, 32(50), 18087-18100.
- Keers, R., Coleman, J. R., Lester, K. J., Roberts, S., Breen, G., Thastum, M., . . . Meiser-Stedman, R. (2016). A Genome-Wide Test of the Differential Susceptibility Hypothesis Reveals a Genetic Predictor of Differential Response to Psychological Treatments for Child Anxiety Disorders. *Psychotherapy and Psychosomatics*, 85(3), 146-158.

- Kovas, Y., Haworth, C. M., Dale, P. S., Plomin, R., Weinberg, R. A., Thomson, J. M., & Fischer, K. W. (2007). The genetic and environmental origins of learning abilities and disabilities in the early school years. *Monographs of the Society for Research in Child Development*, i-156.
- Lionetti, F., Aron, A., Aron, E. N., Burns, G. L., Jagiellowicz, J., & Pluess, M. (2018). Dandelions, tulips and orchids: evidence for the existence of low-sensitive, medium-sensitive and high-sensitive individuals. *Translational psychiatry*, 8(1), 24. doi:10.1038/s41398-017-0090-6
- Meyer-Lindenberg, A., Buckholtz, J. W., Kolachana, B., Hariri, A. R., Pezawas, L., Blasi, G., . . . Weinberger, D. R. (2006). Neural mechanisms of genetic risk for impulsivity and violence in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 103(16), 6269-6274.
- Moore, S. R., & Depue, R. A. (2016). Neurobehavioral foundation of environmental reactivity. *Psychological Bulletin*, 142(2), 107-164. doi:10.1037/bul0000028
- Munafò, M. R., Brown, S. M., & Hariri, A. R. (2008). Serotonin transporter (5-HTTLPR) genotype and amygdala activation: a meta-analysis. *Biological Psychiatry*, 63(9), 852-857.
- Pluess, M. (2015). Individual Differences in Environmental Sensitivity. *Child Development Perspectives*, 9(3), 138-143. doi:10.1111/cdep.12120
- Pluess, M., Assary, E., Lionetti, F., Lester, K. J., Krapohl, E., Aron, E. N., & Aron, A. (2018). Environmental sensitivity in children: Development of the Highly Sensitive Child Scale and identification of sensitivity groups. *Developmental Psychology*, 54(1), 51-70. doi:10.1037/dev0000406

- Pluess, M., & Belsky, J. (2011). Prenatal programming of postnatal plasticity? *Development and Psychopathology*, *23*(1), 29-38.
doi:10.1017/S0954579410000623
- Pluess, M., & Belsky, J. (2013). Vantage sensitivity: Individual differences in response to positive experiences. *Psychological Bulletin*, *139*(4), 901-916.
doi:10.1037/a0030196
- Pluess, M., Stevens, S., & Belsky, J. (2013). Differential Susceptibility: Developmental and Evolutionary Mechanisms of Gene-Environment Interactions. In M. Legerstee, D. W. Haley, & M. H. Bornstein (Eds.), *The Infant Mind: Origins of the Social Brain* (pp. 77-96). New York: Guilford.
- Preacher, K. J., Curran, P. J., & Bauer, D. J. (2006). Computational Tools for Probing Interactions in Multiple Linear Regressions, Multilevel Modeling, and Latent Curve Analysis. *Journal of Educational and Behavioral Statistics*, *31*(4), 437-448.
- Rasch, B., Spalek, K., Buholzer, S., Luechinger, R., Boesiger, P., de Quervain, D. J., & Papassotiropoulos, A. (2010). Aversive stimuli lead to differential amygdala activation and connectivity patterns depending on catechol-O-methyltransferase Val158Met genotype. *Neuroimage*, *52*(4), 1712-1719.
doi:10.1016/j.neuroimage.2010.05.054
- Rijsdijk, F. V., Viding, E., De Brito, S., Forgiarini, M., Mechelli, A., Jones, A. P., & McCrory, E. (2010). Heritable variations in gray matter concentration as a potential endophenotype for psychopathic traits. *Archives of General Psychiatry*, *67*(4), 406-413.
- Roisman, G. I., Newman, D. A., Fraley, R. C., Haltigan, J. D., Groh, A. M., & Haydon, K. C. (2012). Distinguishing differential susceptibility from

diathesis-stress: Recommendations for evaluating interaction effects.

Development and Psychopathology, 24(2), 389-409.

doi:10.1017/S0954579412000065

Ruigrok, A. N., Salimi-Khorshidi, G., Lai, M. C., Baron-Cohen, S., Lombardo, M. V., Tait, R. J., & Suckling, J. (2014). A meta-analysis of sex differences in human brain structure. *Neuroscience and Biobehavioral Reviews*, 39, 34-50.

doi:10.1016/j.neubiorev.2013.12.004

Schriber, R. A., Anbari, Z., Robins, R. W., Conger, R. D., Hastings, P. D., & Guyer, A. E. (2017). Hippocampal Volume as an Amplifier of the Effect of Social Context on Adolescent Depression. *Clinical Psychological Science*, 2167702617699277.

Schwartz, C. E., Kunwar, P. S., Greve, D. N., Kagan, J., Snidman, N. C., & Bloch, R. B. (2012). A phenotype of early infancy predicts reactivity of the amygdala in male adults. *Molecular Psychiatry*, 17(10), 1042-1050.

doi:10.1038/mp.2011.96

Schwartz, C. E., Wright, C. I., Shin, L. M., Kagan, J., & Rauch, S. L. (2003). Inhibited and uninhibited infants "grown up": adult amygdalar response to novelty. *Science*, 300(5627), 1952-1953. doi:10.1126/science.1083703

Sergerie, K., Chochol, C., & Armony, J. L. (2008). The role of the amygdala in emotional processing: a quantitative meta-analysis of functional neuroimaging studies. *Neuroscience and Biobehavioral Reviews*, 32(4), 811-830.

Slagt, M., Dubas, J. S., Dekovic, M., & van Aken, M. A. (2016). Differences in Sensitivity to Parenting Depending on Child Temperament: A Meta-Analysis. *Psychological Bulletin*. doi:10.1037/bul0000061

Whittle, S., Yap, M. B. H., Sheeber, L., Dudgeon, P., Yucel, M., Pantelis, C., . . .

Allen, N. B. (2011). Hippocampal volume and sensitivity to maternal aggressive behavior: A prospective study of adolescent depressive symptoms.

Development and Psychopathology, 23(1), 115-129.

doi:10.1017/s0954579410000684

Whitwell, J. L. (2009). Voxel-based morphometry: an automated technique for

assessing structural changes in the brain. *Journal of Neuroscience*, 29(31),

9661-9664. doi:10.1523/JNEUROSCI.2160-09.2009

Wolf, M., van Doorn, G. S., & Weissing, F. J. (2008). Evolutionary emergence of

responsive and unresponsive personalities. *Proceedings of the National*

Academy of Sciences of the United States of America, 105(41), 15825-15830.

doi:10.1073/pnas.0805473105

Yap, M. B., Whittle, S., Yucel, M., Sheeber, L., Pantelis, C., Simmons, J. G., & Allen,

N. B. (2008). Interaction of parenting experiences and brain structure in the

prediction of depressive symptoms in adolescents. *Archives of General*

Psychiatry, 65(12), 1377-1385. doi:10.1001/archpsyc.65.12.1377

Zuckerman, M. (1999). *Vulnerability to psychopathology: A biosocial model*.

Washington: American Psychological Association.

Table 1*Demographic Characteristics of the Sample (N = 62)*

Variables	N (%)
Age at imaging (years)	$M = 11.61, SD = .81$ (Range: 10.17 - 13.17)
Age at outcome (years)	$M = 12.09, SD = .29$ (Range: 11.54 – 12.73)
Zygoty	
MZ	20 (32.3%)
DZ	42 (67.7%)
Ethnicity	
White	57 (91.9%)
Other	5 (8.1%)
Annual Household Income	
< £17,499	7 (11.3%)
£17,500 - £49,999	41 (66.1%)
> £50,000	11 (17.7%)
Brain Volume	
Amygdala Left Volume	$M = .79, SD = .07$
Amygdala Right Volume	$M = .59, SD = .05$
Hippocampus Left Volume	$M = .73, SD = .04$
Hippocampus Right Volume	$M = .64, SD = .03$

Environment Quality¹ $M = 13.88, SD = 1.67$

Total Problem Behaviors (SDQ) $M = 6.50, SD = 5.57$

Note. ¹ Higher scores reflect higher quality of the environment. The Environmental Quality score ranges from 8 = lowest quality to 18 = highest quality.

Table 2*Unadjusted Associations between Variables (N = 62)*

Variables	1	2	3	4	5
1 Amygdala Left Volume	—				
2 Amygdala Right Volume	.66**	—			
3 Hippocampus Left Volume	.52**	.46**	—		
4 Hippocampus Right Volume	.46**	.58**	.77**	—	
5 Environmental Quality ¹	-.10	-.13	.12	.08	—
6 Total Problems	-.21	-.07	.06	.12	-.14

Note. ¹ Higher scores reflect higher quality of the environment. * $p < .05$. ** $p < .01$.

Table 3*Summary of Hierarchical Linear Models Predicting Total Problems (N = 62)*

Predictor Variables	Total Problems			
	Model AL	Model AR	Model HL	Model HR
Step 1				
Amygdala Left Volume	-13.10	—	—	—
Amygdala Right Volume	—	-6.85	—	—
Hippocampus Left Volume	—	—	8.46	—
Hippocampus Right Volume	—	—	—	30.78
Environmental Quality	-.49	-.45	-.44	-.45
Step 2				
Amygdala Left X Environment	-16.10*	—	—	—
Amygdala Right X Environment	—	-12.32	—	—
Hippocampus Left X Environment	—	—	3.48	—
Hippocampus Right X Environment	—	—	—	-11.73

Note. The displayed coefficients of variables at step 1 represent the values before inclusion of interaction term at step 2; AL = Amygdala Left; AR = Amygdala Right; HL = Hippocampus Left; HR = Hippocampus Right; * $p < .05$. ** $p < .01$.

Figure Legends

Figure 1.

Illustration of three models of Environmental Sensitivity. *Diathesis-Stress* describes individual differences in response to exclusively negative influences whereas *Vantage Sensitivity* refers to variability regarding positive influences only. *Differential Susceptibility* represents the combination of diathesis-stress and vantage sensitivity with heightened sensitivity to both negative and positive experiences (based on Figure 1 in Pluess, 2015).

Figure 2.

Graphic illustration of the selected brain regions used to generate the amygdala and hippocampus volumes. Left and right amygdala are colored red whereas left and right hippocampus are indicated as blue.

Figure 3.

Simple slopes and scatter plot of the significant interaction between left amygdala and environmental quality predicting total behavior problems. The hierarchical linear models were run with continuous brain volumes but for the follow-up analysis the sample was divided by medium split into small amygdala (triangles in scatter plot) and large amygdala (dots in scatter plot) (some of the triangles and dots are overlapping). Shaded areas reflect regions of significance. In these regions the association between amygdala left volume and behavior problems is significant.